



NURSING ASSESSMENTS:

The following provides a guideline for clinical assessment. Assessments must be done at regular intervals and are used to drive treatment decisions.

- **Assessment of risk or contributing factors** associated with skin breakdown should be determined from the patient's history. A summary of risk factors and types of impaired skin integrity follows:

Problem	Risk Factors	
	Local	Systemic
Stripping/Shearing Injury	Mechanical Trauma	Elderly
	Ischemia	Immobility
	Dry Tissue	Malnutrition
Perineal Skin Compromise	Incontinence	Recent antibiotic use
	Inadequate hygiene	Elderly
	Mechanical trauma	Malnutrition
Arterial Ulcer	Mechanical trauma	Peripheral vascular disease
		Smoking
Neuropathic Ulcer	Mechanical trauma	Diabetes
		Spinal cord injury
Pressure Ulcer	Moisture	Malnutrition
	Mechanical trauma	Immobility
		Decreased activity
		Sensory perception deficits
Venous Ulcer	Edema	History of deep vein thrombosis
	Cellulitis	Previous leg ulceration
	Mechanical trauma	Obesity

- **General assessments:** include patient's current health status, disease processes, age, activity level, nutrition and medications.
- **Assessment of the skin:** color; temperature; sensation (e.g. pain, itching); hydration (e.g. dry, cracked, moist); tissue consistency (e.g. firm, boggy); thin skin; presence of edema; induration; changes in any of the above; and, presence of healed ulcer or incision.
- **Assessment of nutrition, hydration, current skin care, patient/caregiver level of understanding, compliance in care, and learning style.**

GENERAL NURSING INTERVENTIONS

- **Provide systematic skin inspection** at least daily based upon characteristics listed above.
- **Implement prevention protocols** based on the potential for the following:
 - Stripping/Shearing Injury* - proper turning, positioning techniques, careful selection and removal of adhesives, use of alternatives to tape.
 - Perineal Skin Compromise* - cleanse and protect tissue at frequent intervals, gently cleanse skin, frequent use of a moisturizer or barrier is recommended with incontinence, appropriate use of incontinence containment products (e.g. fecal collectors, external urinary collection pouches, external urinary catheters) is recommended.
 - Arterial Ulcer* - consult with physician regarding ischemia and planned treatment, protect extremity from trauma, cleanse, moisturize and protect intact skin, avoid foot soaks.
 - Neuropathic Ulcer* - consult with physician if ischemia exists, protect from injury with orthotics or other appropriate footwear, cleanse, moisturize and protect intact skin, avoid foot soaks.
 - Pressure Ulcer* - proper positioning, turning, patient support surface and/or wheelchair seat are essential, cleanse, moisturize and protect intact skin, avoid massage of bony prominences, orthotic devices.
 - Venous Ulcer* - compression therapy, leg elevation, exercise (e.g. walking), weight management as needed, cleanse, moisturize and protect intact skin.
- **Provide adequate nutritional intake and hydration.** Education plans should include compliance with care, medications and preventive approaches (e.g. smoking cessation, weight control).
- **Provide education:** patient, family and caregiver.
- **Document** assessments and interventions.
- **Reassess** at regular intervals per agency protocol.

BACKGROUND INFORMATION:

The **skin** is the largest organ of the body and, as such, comprises a surface area which is subjected to external injury from mechanical forces (e.g. pressure, friction, shear, stripping), chemical exposure (e.g. urine, stool, wound exudate, solutions for skin and wound care), radiation (e.g. ultraviolet, therapeutic) and potential pathogens (e.g. fungi, bacteria). Provision of a barrier is one of the skin's primary functions. Additionally, it prevents dehydration through evaporative water loss, maintains thermoregulation, synthesizes vitamin D and provides sensory feedback.

Anatomically, the skin is composed of two layers: the epidermis and the dermis.

The **epidermis**, which is the outermost layer of the skin, is characterized as follows:

- avascular
- varies in thickness (depending on body location)
- a dry structure which sheds cells and replaces itself every 4-6 weeks
- approximately the thickness of a piece of plastic wrap

The **dermis** is located directly beneath the epidermis and is characterized as follows:

- provides strength and structural support through a vascular network of connective tissues
- contains blood vessels, nerves, hair, nails, sebaceous glands and sweat glands
- is thicker than the epidermis

Below the dermis, is the **subcutaneous tissue** which is composed of major vessels, lymphatics, fat and connective tissue. This area provides insulation and nutritional support for the skin. Located below the subcutaneous tissue are **fascia, muscles, tendons and bone**. All layers of tissue below the epidermis are moist. Therefore, moisture retentive wound care treatments are usually indicated in order to maintain cell life and proliferation.

The **skin changes as we age**. The number of sweat glands declines and the epithelial and fatty layers of the subcutaneous tissue become thinner. As this padding is lost, a higher risk of skin breakdown secondary to pressure, friction, stripping and shearing exists. Itching and dry skin are also common complaints. Disease states, dehydration, malnutrition, medications and immobilization may further affect the skin and, when a wound is present, may impair healing.

Variations in **skin color** based on ethnic background can lead to a missed diagnosis of early compromise. Although the epidermal outer layer in black patients is the same thickness as that of lighter skinned individuals, there are a greater numbers of cells which are arranged in a more compact fashion. This results in a more effective barrier to chemical and mechanical insults. However, the dark pigmentation also makes assessment of early injury and treatment more difficult to detect. In black patients, an "ashy" appearance of the skin, warmth, tightness or firmness under the skin are signs of early compromise.

An in-depth discussion of the skin is beyond the scope of this material. However, knowledge of optimal conditions for healthy skin is important to understanding the rationale for prevention techniques.

The algorithm on the reverse side provides a general path of decision-making for assessment, prevention and management of early skin breakdown. Below is detailed information designed to assist health care providers. This tool should be used along with the consultative services of a skin and wound care specialist such as a WOC nurse, physical therapist, clinical nurse specialist with expertise in skin care or a physician when indicated.



Intact Skin

Intact skin is without visible evidence of injury or where there is a healed wound which has epithelialized and maintains closure.

Goals of Care: protect and maintain intact skin.

Wound and Skin Care Objectives: cleanse and moisturize intact skin.



Compromised Skin

Compromised skin is tissue exposed to potential injury or tissue that is in a weakened condition (e.g. dry, thin).

Goals of Care: maintain intact skin and improve tissue tolerance.

Wound and Skin Care Objectives: cleanse and moisturize skin and protect tissue.

REFERENCES

- Fiers SA. Breaking the cycle: the etiology of incontinence dermatitis and evaluating and using skin care products. *Ostomy/Wound Management* 1996 42(3) 32-43.
- Hess, Cathy Thomas. *Nurse's Clinical Guide to Wound Care*. Springhouse, PA, Springhouse Corporation, 1995.
- Krasner D, Kane D (eds). *Chronic Wound Care, A Clinical Source Book for Healthcare Professionals, Second Edition*, Wayne, Pennsylvania, Health Management Publications, Inc. 1997.
- Maklebust J, Sieggreen. *Pressure Ulcers: Guidelines for Prevention and Nursing Management, Second Edition*. Springhouse, PA, Springhouse Corporation, 1996.
- Morison M, Moffatt C: *A Colour Guide to the Assessment and Management of Leg Ulcers, Second Edition*, London, Mosby 1994.
- Panel for the Prediction and Prevention of Pressure Ulcers in Adults. *Pressure Ulcers in Adults: Prediction and Prevention. Clinical Practice Guideline, No. 3. AHCPR Publication No. 92-0047*. Rockville, MD: Agency for Health Care Policy and Research, Public Health Service, U.S. Department of Health and Human Services, May 1992.
- Wound, Ostomy and Continence Nurses Society. *Guidelines For Management: Caring For A Patient With Fecal Incontinence*. Laguna Beach, CA 1994.
- Wound, Ostomy and Continence Nurses Society. *Standards of Care: Patient With Urinary Incontinence*. Laguna Beach, CA 1992.
- Wysocki AB. A review of the skin and its appendages. *Advances in Wound Care* 1995 8(2) 53-70.

Skin at Risk

- Restore Clean 'N Moist
- Hollister Skin Cleanser
- Restore Barrier Creme
- Hollister Skin Conditioning Creme
- Hollister Moisture Barrier Ointment
- Hollister Skin Gel
- Restore Extra Thin Hydrocolloid
- EpiFlex Heel and Elbow Protectors

*Developed in collaboration with Bonnie Sue Rolstad, RN, BA, CWOON
Bryant Rolstad Consultants, LLC, St. Paul, MN
Photography courtesy of Ms. Rolstad*



2000 Hollister Drive
Libertyville, IL 60048 USA
©1999 Hollister Incorporated
All Rights Reserved.
Printed in USA

Hollister and design, Hollister, Restore, Restore Cx, Restore CalciCare and EpiFlex are trademarks of Hollister Incorporated.