Address intubated patient risks with confidence.

Every patient dependent on a ventilator for respiration is vulnerable. Ventilator-associated risks can result in increased length of stay and increased hospital costs.¹

RISKS of intubated patients	CLINICAL BENEFITS of the AnchorFast oral endotracheal tube fastener
VENTILATOR-ASSOCIATED EVENTS such as pneumonia, due to limited access to oral care.	EASES ACCESS TO THE ORAL CAVITY to optimize oral care. Findings suggest implementation of a comprehensive oral and dental care protocol can help reduce incidence of ventilator-associated pneumonia. ²
ORAL PRESSURE ULCERS from the tube resting too long in one place.	HELPS PREVENT FORMATION OF LIP ULCERS as part of a multifaceted approach. ^{3, 4}
UNEXPECTED TUBE OCCLUSIONS caused by the patient.	HELPS PREVENT TUBE OCCLUSION (AnchorFast Guard only) with integrated tube protection sleeve.

AnchorFast Oral Endotracheal Tube Fastener

•••••

 STOCK NO.	PRO
9800	Anc Oral
9799	Anc Oral
9788	Anc Rep

Hollister Incorporated Libertyville, IL USA	¹ Res on 31(
For orders only: 1.800.323.4060	² Ga ora (20
For detailed clinical questions concerning	³ Mil to F Res
our products, call: 1.888.740.8999	⁴ Ver the 17-
Distributed in Canada by:	
Hollister Limited	
Aurora, Ontario	CAL
1.800.263.7400	a ph
	be s
www.hollister.com	the p Prec



T



Restrepo M, et al. "Economic Burden of Ventilator-Associated Pneumonia Based on Total Resource Utilization." Infection Control and Hospital Epidemiology, 2010; 31(5):509-515.

Carcia R, et al. "Reducing ventilator-associated pneumonia through advanced ral-dental care: a 48-month study." American Journal of Critical Care 18.6 2009): 523-32.


filler, Kenneth, et al. "A Utilization of AnchorFast Oral Endotracheal Tube Fastener b Reduce the Incidence of Lip Ulcers." Presented at The 55th International tespiratory Congress. 5-8 December 2009, San Antonio, Texas USA.

eneman, Wade, et al. "An Alternative Approach to Prevent Lip Pressure Ulcers in ne Intubated Patient." Presented at the Symposium on Advanced Wound Care. 7-20 April 2010, Orlando, Florida USA.

AUTION: Federal (USA) Law restricts this device to sale by or on the order of obysician. Prior to use of the AnchorFast Guard oral endotracheal tube fastener, a sure to read the entire product Instructions for Use package insert supplied with a product for device intended Use, Description, Contraindications, Warnings, ecautions, Adverse Events and Instructions for Use.

Hollister and logo, AnchorFast, and AnchorFast Guard are trademarks of Hollister Incorporated. © 2016 Hollister Incorporated. Printed in USA 922960-0116

When your choice is too important to take a chance.







History, quality, and innovation.

Used at over 3,700 hospitals in the United States

In thousands of hospitals around the world, experts and professionals depend on our AnchorFast products to help improve patient care.

20 years of demonstrated quality

Hollister has a long history with endotracheal tube securement devices. Clinicians depend on its superior performance and quality, making AnchorFast products overwhelmingly the best-selling endotracheal tube securement devices on the market.

Ongoing innovation driven by customer feedback

Innovation happens when we listen to the people who use our products. What they say matters to us. It's why AnchorFast products continue to evolve to meet your needs.


"It's important to look for vendors who have an interest beyond just the transaction. Hollister is more than a supplier, they're a partner to respiratory care practitioners in our continuous quest for advancement in patient care." — Respiratory Care Therapist

Adjustable neck strap and is replaceable.



AnchorFast Oral Endotracheal Tube Fastener

The **Anchor**Fast oral endotracheal tube fastener combines a number of unique features that make it the right choice.



AnchorFast Guard

Oral Endotracheal Tube Fastener

With the addition of integrated tube protection to our trusted securement system, the AnchorFast Guard tube fastener can help bring a new level of confidence to the ICU.